

Your contact at PhytoLab:
Reference Substances
Tel.: +49 9163 88-395
ref-substances@phytolab.de
<https://phyproof.phytolab.com>

Certificate of analysis

Article:	89514 Campesterol
Certificate # / Lot Number:	70209
Material batch:	20366
Sample-ID:	38371
End of analysis:	04/2022
Expiry date:	08/2025

Test	Unit	Specified value	Testresult
Appearance, SOP 100005		powder	conform
Color, SOP 100006		white	conform
Identification (1H-NMR-spectroscopy), (outsourced), SOP 206010		conform	conform
Identification (13C-NMR-spectroscopy), (outsourced), SOP 206020		conform	conform
Identification (IR-spectroscopy, Ph.Eur. 10.3, 2.2.24 / USP43 NF37 <197>), SOP 206000		conform	conform
Water content, (micro determination, coulometric titration), Ph.Eur. 10.0., 2.5.32, SOP 304291 Vers. 2018-01: Mean value	%		0.2
Campesterol (GC), method 1 (% AU), SOP 441660	%	≥ 90.00	96.81
Inorganic impurities, (ICP-MS), for reference substances, SOP 811701: Calcium	%		<0.1
Potassium	%		<0.1

Certificate of analysis

Article: 89514 Campesterol
 Material batch: 20366

Test	Unit	Specified value	Testresult
Magnesium	%		<0.1
Sulfur	%		<1.0
Sodium	%		<0.1
Phosphorus	%		<0.1
Aluminium	%		<0.1
Residual solvents, (headspace-GC), SOP 805765:	%		
Residual solvents (LOQ: 0.050)			<0.050
Content, SOP 890011, calculated in (%): (100 - water - residual solvents - inorganic impurities) x chromatographic purity / 100 x factor (SOP 206070) For detailed information refer to attached data sheet! Campesterol	%		53
Dihydrobrassicasterol	%		43
Sum of campesterol and dihydrobrassicasterol	%		96

This PhytoLab phyproof© reference standard is by definition a primary reference standard and does not need to be qualified against any other reference standard. The identity of the reference standard has been substantiated by at least two independent analytical methods such as IR, NMR, UV or MS analysis. A mass balance approach, which takes chromatographic purity into account, as well as the contents of water, residual solvents, inorganic impurities, and the counter ion (if the reference standard is present as a salt) is applied in the calculation of the absolute purity as given in this COA (see description of SOP 8900XX).

The absolute purity value (and not just the chromatographic purity result obtained by means of HPLC or GC) must be used in all quantitative calculations as the chromatographic techniques do not yet account for water, residual solvents and inorganic impurities.

Certificate of analysis

Article: 89514 Campesterol
Material batch: 20366

The reference substance campesterol always contains a certain percentage of its epimer dihydrobrassicasterol. The two compounds can hardly be separated chromatographically. Therefore, the purity of > 90 % as determined by GC actually describes the sum of both substances. However, the ratio between campesterol and dihydrobrassicasterol can be determined by the evaluation of the intensities of the respective carbon signals in the ¹³C-NMR spectra (SOP 206070, 206080). The resulting factors are included in the calculation of the absolute purity value according to SOP 890011 - shown here are the absolute purity values for campesterol, dihydrobrassicasterol and the sum of both sterols. Please note that due to possible different ratios of epimers in the investigated sample and the reference substance, no conclusions can be drawn regarding the content of the individual substances campesterol and dihydrobrassicasterol. However, using the total sterol content (sum of campesterol and dihydrobrassicasterol) of the reference substance, the total sterol content in the tested samples can be calculated. Please also refer to the enclosed data sheet for further information.

Vestenbergsreuth, 02/Aug/2023

Lisa Fuhrmann

QC Reference Substances

This is a computer print and valid without signature. A signed certificate of analysis can be taken on request.

Certificate of analysis

Article: 89514 Campesterol
Material batch: 20366

Further information:

Shelf life/stability: The stated [expiry](#) date applies when the reference substance is stored in the original unopened container within the specified temperature range. PhytoLab does not guarantee the stability of the reference substance once the vial has been opened.

Long-term storage and handling: The reference standard should be stored in the original unopened vial, protected against light and humidity in an airtight container, within the temperature range given on the label and accompanying data sheet. If stored below room temperature, the vial should be warmed up to room temperature in a desiccator before it is opened in order to avoid condensation of humidity. The user assumes responsibility for deciding how previously opened reference standard vials should be used and the user must ensure that the contents of opened vials are still suitable for their intended use.

Exact weight: the exact weight of each vial is given on the label of the inner vial to two decimal places. This information may be used to produce stock solutions of a known concentration without having to weigh in the reference substance again. If used for this purpose, the content of the vial must be quantitatively transferred to a volumetric flask and filled up to the required level. Please note that PhytoLab is unable to guarantee the stability of the reference standard in solution.

Intended use: this reference standard is solely intended for laboratory analytical purposes, research & development, and scientific teaching and training purposes. It may not be used for any other purpose and particularly not for use in, or the production of, food, animal feed, human or veterinary drugs, cosmetics, medicinal products or diagnostic agents, including in-vitro diagnostic agents. PhytoLab is unable to guarantee the suitability of this reference standard for any particular application other than its qualitative and quantitative use in chromatography and identification testing.

Further information about this reference standard can be found on the accompanying data sheet or in our webshop. Spectral and chromatographic data, and a description of the applied chromatographic method, are provided in the attachments to this COA. A detailed explanation of all data given on the COA can be found in the guide that is available from the download area in our webshop, where you can also download all of the safety data sheets.


Product Data Sheet

Campesterol

Product #: 89514

Physicochemical Data

CAS #:	474-62-4
Molecular formula:	C ₂₈ H ₄₈ O
Molecular weight [g/mol]:	400.68
Synonyms	Campesterin; 24 α -Methylcholesterol
Substance class:	Isoprenoids
Subgroup 1:	Steroid-type
Subgroup 2:	Sterols
Solubility:	soluble in methanol; practically insoluble in water Please note that this solubility information is based on in-house experience or taken from published data. It is not meant to guarantee solubility up to a specific concentration, nor does it guarantee stability of the reference substance in solution.



Additional Information

Please note: The reference substance campesterol always contains a certain percentage of its epimer dihydrobrassicasterol. The epimers cannot be separated chromatographically. However, the ratios between campesterol and dihydrobrassicasterol can be determined by the evaluation of the intensity of the respective carbon signals in the ¹³C-NMR spectra. From this ratio a factor is derived which is included in the calculation of the absolute purity for campesterol, dihydrobrassicasterol and the sum of both sterols.

Source: botanical origin

Long-term storage conditions: 2-8 °C

Manufacturer: Phytolab GmbH & Co.KG
Dutendorfer Straße 5-7
91487 Vestenbergsgreuth
Germany

Tel.: +49 9163 88-395
Fax: +49 9163 88-456
Mail: ref-substances@phytolab.de
Shop: <https://phyproof.phytolab.com>



Supplements


Campesterol
Product # 89514

Batch # 20366

batch 20366 corresponds to batch 19664 which was analysed completely

Identity tests:

IR spectrum




C:\Users\Public\Documents\Bruker\OPUS_ProtectedPool\MEAS\89514_Campesterol_19664.0




NMR spectra

¹H-NMR (CDCl₃; 500 MHz)




¹³C-NMR (CDCl₃; 125 MHz)





Chromatographic purity:



Peak #	RetTime [min]	Sig	Type	Area [pA*s]	Area %	Name
1	39.535	1	BB	299.56451	96.84323	Camosterol
2	40.718	1	MM	9.76480	3.15677	?

Analytical conditions

Column: DB-1; L: 30 m; Id: 0.25 mm; FT: 0.25 μ m
Carrier Gas: Nitrogen 55 kPa constant pressure
temperature gradient: 150°C / 3 min isoth 30°C/min. – 260 °C – 1°C/min to 300°C / 15.0 min isoth.
Injection Volume: 1 μ l, Split ratio:1:30
Injector Temperature: 230 °C
Sample concentration: approx. 1.0 mg/1 ml
Sample preparation: dissolved in dichloromethane
Detection: FID 280 °C (O₂: 400 ml/min. H₂: 40 ml/min, Make-up N₂: 35 ml/min)
Special note: -

Please note: Values on the certificate of analysis may vary as these are average values of at least three injections while above chromatogram and report is only one example. Non-integrated peaks originate from the blank injection.